




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷: A61K	A2	(11) International Publication Number: WO 00/24362 (43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/24486 (22) International Filing Date: 20 October 1999 (20.10.99) (30) Priority Data: 60/105,850 27 October 1998 (27.10.98) US 09/273,766 22 March 1999 (22.03.99) US (71) Applicant: VIRGINIA COMMONWEALTH UNIVERSITY [US/US]; 1101 E. Marshall Street, Richmond, VA 23298 (US). (72) Inventors: PEART, Joanne; Apartment #1332, 11808 Chase Wellesley Drive, Richmond, VA 23233 (US). BYRON, Peter; 8101 Lower Ralston Court, Richmond, VA 23229 (US). LICHTMAN, Aron; 2306 Stonemill Court, Richmond, VA 23233 (US). MARTIN, Billy; 2406 St. Regis Drive, Richmond, VA 23233 (US). (74) Agent: WHITMAN, Michael, E.; Whitham, Curtis & Whitham, Suite 900, 11800 Sunrise Valley Drive, Reston, VA 20191 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: Δ^9 TETRAHYDROCANNABINOL (Δ^9 THC) SOLUTION METERED DOSE INHALERS AND METHODS OF USE		
(57) Abstract The present invention provides therapeutic formulations for solutions of Δ^9 -tetrahydrocannabinol (Δ^9 THC) to be delivered by metered dose inhalers. The formulations, which utilize non-CFC propellants, provide a stable aerosol-deliverable source of Δ^9 THC for the treatment of various medical conditions, such as: nausea and vomiting associated with chemotherapy; muscle spasticity; pain; anorexia associated with AIDS wasting syndrome; epilepsy; glaucoma; bronchial asthma; and mood disorders.		



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Δ^9 TETRAHYDROCANNABINOL (Δ^9 THC) SOLUTION METERED DOSE INHALERS AND METHODS OF USE

DESCRIPTION

BACKGROUND OF THE INVENTION

Field of the Invention

The invention is generally related to the therapeutic use of Δ^9 Tetrahydrocannabinol (Δ^9 THC). In particular, the invention provides a metered dose inhaler (MDI) for the aerosol administration of Δ^9 THC to patients suffering from nausea and vomiting associated with cancer chemotherapy, muscle spasticity, pain, anorexia associated with AIDS wasting syndrome, epilepsy, glaucoma, bronchial asthma, mood disorders, and the like.

Background Description

"Medical Marijuana" is a timely and controversial subject that is currently receiving widespread public attention. While marijuana is usually thought of as an illegal "recreational" drug, it also has a long history as a medicine. In 1997, the National Institutes of Health (NIH) released a review of the scientific data concerning potential therapeutic uses for marijuana. In that review, the NIH found that marijuana may indeed have beneficial medicinal effects and recommended that researchers develop alternative dosage forms for the drug, such as a "smoke free" inhaled delivery system (1). Table 1 summarizes the findings of several studies (references 2-18) that have documented therapeutically beneficial medicinal uses of the major active component of marijuana, Δ^9 tetrahydrocannabinol (Δ^9 THC).

TABLE 1. The Use of Δ^9THC for the Treatment of Assorted Clinical Conditions			
Condition and Number of Patients	Administration Route and Dose	Findings	Reference
AIDS-associated anorexia and cachexia; 94 patients; 12 months	Oral placebo, 2.5 mg THC once or twice daily increasing to 20 mg daily	Long term THC treatment was well-tolerated; THC improved appetite and only tended to increase weight compared to controls	Beal et al., 1997
AIDS-associated anorexia and cachexia; 139 patients; 42 days	Oral placebo or 2.5 mg THC twice daily	57% and 69% of vehicle and THC patients were evaluable for efficacy. Appetite increased 38% over baseline for THC group compared to only 8% for the placebo group. THC also decreased nausea. No significant changes were found between the groups for weight change.	Beal et al., 1995
Nausea and emesis due to cancer chemotherapy; 36 patients who had experienced severe nausea and vomiting that was refractory to prochlorperazine or thiethylperazine	Oral THC, 15mg/m ²	Reduction in chemotherapy-induced nausea and vomiting in 64% of patients given THC compared to prochlorperazine; side effects included dysphoria; authors recommend initial THC dose of 5 mg/m ²	McCabe et al., 1988

TABLE 1. The Use of Δ^9 THC for the Treatment of Assorted Clinical Conditions

Nausea and emesis due to cancer chemotherapy; 53 patients which were refractory to other antiemetics	Oral 5 or 15 mg/m ² THC four times per day	72% of patients exhibited a THC-induced partial or complete blockade of vomiting	Lucas and Laszlo, 1980
Nausea and emesis due to cancer chemotherapy; 84 patients	Oral 10 mg/m ² THC of prochlorperazine	THC more effective than prochlorperazine	Sallan et al., 1980
Nausea and emesis due to cancer chemotherapy; 116 patients	Oral 15 mg THC, 10mg prochlorperazine or placebo	Equal antiemetic effects between THC and prochlorperazine. effects of each greater than placebo; considerably more CNS side effects with THC than prochlorperazine	Frytak et al., 1979
Nausea and emesis due to cancer chemotherapy; 15 patients	Oral placebo or 10 mg/m ² THC every 3 hours for a total of 5 doses, THC (17 mg) laced cigarettes of placebo were given if vomiting occurred	93% patients had a reduction in nausea and vomiting, 53% had an excellent response, 40% had a fair response; plasma THC levels 7.1±6.9 (mean ± SD) ng/ml. Side effects included sedation, tachycardia, few other side effects	Chang et al., 1979
Pain due to advanced cancer; 10 patients	Oral placebo and 5, 10, 15 or 20mg THC	Pain relief, elevated mood, appetite stimulation, drowsiness, slurred speech, mental clouding	Noyes, et al, 1975

TABLE 1. The Use of Δ^9 THC for the Treatment of Assorted Clinical Conditions

Pain due to advanced cancer; 34 patients	Placebo, 10 and 20mg THC, and 60 and 120 codeine	THC produced a similar degree of analgesia, with greater potency than codeine. THC CNS side effects included sedation, mental clouding, ataxia, and disorientation	Noyes et al. 1975
Spasticity related to multiple sclerosis; 2 patients	Oral 10 or 15mg THC; rectal dose of 5 or 10mg THC	Improvement in passive mobility and walking ability	Brenneisen et al., 1996
Spasticity related to multiple sclerosis; 13 patients	Oral 2.5 to 15mg THC once or twice daily or placebo	Significant subjective improvement in spasticity at 7.5mg THC and higher, no significant improvement in objective measurements	Ungerleider et al., 1987
Spasticity related to multiple sclerosis; 8 patients, single blind	Oral 5 to 15mg THC	5 of 8 patients had mild subjective improvement in tremor. 2 of 8 patients had both objective and subjective improvement	Clifford, 1983
Spasticity related to multiple sclerosis; 9 patients	Placebo, or 5 or 10mg THC	Decrease in spasticity compared to placebo treatment, minimal side effects	Petro and Ellenberger, 1981
Spasticity and pain due to spinal cord injury; 1 patient	Oral placebo, THC (5 mg), or codeine (50 mg)	THC and codeine had analgesic effect compared to the placebo treatment. THC had a beneficial effect on spasticity whereas codeine did not	Maurer et al., 1990

TABLE 1. The Use of Δ^9THC for the Treatment of Assorted Clinical Conditions			
Glaucoma, 6 patients	Oral placebo or 5, 10, 15 and 20 mg THC	Pain relief, elevated mood, appetite stimulation, drowsiness, slurred speech, mental clouding	Merritt et al, 1980
Ten subjects with normal intra ocular pressure	Intravenous THC (0.022 or 0.044 mg/kg)	Decreased intra ocular pressure by a mean of 37%	Cooler and Gregg, 1977
Nausea and emesis due to cancer chemotherapy; refractory to other antiemetics	Oral 10 mg/m ² THC or placebo	In 20 courses of THC, 5 resulted in no vomiting, 9 resulted in a reduction of vomiting, 3 resulted in no decrease in vomiting, and 2 were unevaluable. THC was significantly better than placebo in decreasing vomiting.	Sallan et al., 1975

When marijuana is used illegally as a recreational psychoactive drug, the active ingredient Δ^9 THC is usually delivered to the lungs as an impure non-pharmaceutical aerosol in the form of marijuana smoke. Aerosolized Δ^9 THC in the inhaled smoke is absorbed within seconds and delivered to the brain efficiently. Table 2 and references 19-20 describe the pharmacokinetics of the administration of Δ^9 THC. As can be seen, inhalation is the preferred route of delivery for Δ^9 THC. When compared to oral delivery, inhalation provides a more rapid onset of pharmacological action and peak plasma levels. The effects achieved via inhalation are comparable to those achieved when the drug is administered intravenously, but inhalation is a much less invasive technique.

TABLE 2. Pharmacokinetics of Δ^9THC Given Orally, Intravenously or by Smoking					
Route	Dose	% Dose in Plasma	Onset of Pharmacological Action	Peak Plasma Levels	References
Oral, sesame oil in gelatin capsules	2.5, 5, or 10 mg	10 to 20%	0.5 to 1 hour	120-480 min	(PDR, 1995)
Oral, in cookies	20 mg	4 to 12%	120-180 min	60-90 min	(Ohlsson, et al., 1980)
Intravenous, bolus	5 mg	100%	10 min	3 min	(Ohlsson, et al., 1980)
Smoking (THC lost to side stream smoke and pyrolysis)	13 mg	8 to 24%	10 min	3 min	(Ohlsson, et al., 1980)

Currently, the sources of Δ^9 THC for patients who could benefit from the drug are very limited. An oral form of Δ^9 THC (MARINOL) is marketed as a treatment for nausea and vomiting related to cancer chemotherapy, and as an appetite stimulant in patients suffering from AIDS wasting syndrome. In MARINOL, pharmaceutical grade Δ^9 THC is dissolved in sesame oil, encapsulated in gelatin capsules and delivered orally. However, when the drug is taken orally, the absorption is slower and more variable than when inhaled, with an onset of action between 30 minutes and 2 hours (Table 2). Alternatively, some cancer patients do manage to

obtain and smoke marijuana in order to alleviate such conditions as nausea and vomiting due to chemotherapy. This is, however, technically illegal and is thus obviously a less than ideal treatment protocol. There is no currently available pharmaceutically acceptable aerosol form of Δ^9 THC.

It would be advantageous to have available a form of pharmaceutical grade Δ^9 THC that could be administered as an aerosol. This would provide a means for rapid uptake of the drug without resorting to the illegal practice of smoking marijuana. Also, the potential adverse side effects encountered by smoking marijuana would be avoided. Further, an aerosol preparation of pharmaceutically pure Δ^9 THC could be administered in known, controlled dosages.

In 1976, Olsen et al. described a chlorofluorocarbon (CFC) propelled MDI formulation of Δ^9 THC (21). However, Δ^9 THC is known to deteriorate during storage, and the stability of Δ^9 THC in this formulation is suspect. In addition, the ethanol content in this formulation was so high (~23%) as to create an aerosol with droplets too large to be effectively inhaled (22). The Δ^9 THC CFC formulations were tested for use in treating asthma but were shown to be only moderately effective (23, 24). Moreover, CFC propellants have since been banned so that such a formulation is now useless. It would clearly be advantageous to develop a new aerosol formulation in which the Δ^9 THC is stable, the droplets are of a size that can be effectively inhaled, and which utilizes a non-CFC propellant.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide a stable aerosol-dispensable pharmaceutical composition comprising a non-CFC propellant and a pharmaceutically effective concentration of Δ^9 THC. More particularly, it is an object of the present invention to provide a stable aerosol-dispensable pharmaceutical composition comprising a hydrofluoroalkane propellant, (for example, HFA 227 or HFA 134a) and Δ^9 THC. The propellant is present in the range of approximately 78 to 100% by

weight, and more particularly the propellant is present in the range of approximately 85 to 100% by weight. An organic solvent such as ethanol can be used to assist in solubilizing the Δ^9 THC in the propellant but is not required. If a solvent is used, preferably less than 20% by weight will be required, and most preferably less than 15% by weight will be required. The pharmaceutically effective concentration of Δ^9 THC is preferably in the range of 0.05 to 10% by weight, and most preferably in the range of 0.1 to 6% by weight. The pharmaceutical composition of the present invention can be used to treat a variety of medical conditions including nausea and vomiting associated with cancer chemotherapy, muscle spasticity, pain, anorexia associated with AIDS wasting syndrome, anorexia associated with cancer chemotherapy, epilepsy, glaucoma, bronchial asthma, mood disorders, migraine headaches.

DETAILED DESCRIPTION OF THE DRAWINGS

Figure 1. Δ^9 THC MDI characterization summary before and after storage at 40°C and 82% relative humidity (RH).

Figure 2. Generalized schematic drawings of a Δ^9 THC MDI.

DETAILED DESCRIPTION OF A PREFERRED EMBODIMENT OF THE INVENTION

The instant invention provides a series of non-ozone depleting pressurized metered dose inhaler formulations of Δ^9 THC. In preferred embodiments of the invention, the formulations contain the pharmaceutically acceptable, non-ozone depleting hydrofluoroalkane propellants HFA 134a (1,1,1,2-tetrahydrofluoroethane) and HFA 227 (1,1,1,2,3,3,3-heptafluoropropane), or a mixture thereof.

When the propellant is a hydrofluoroalkane, it has been discovered that the propellant may be used with or without a solvent such as ethanol. Higher percentages of solvent generally allow higher levels of dissolution of Δ^9 THC. However, higher

percentages of solvent also cause droplet size to increase. In preferred embodiments of the invention, the range of propellant compositions, as shown in Table 3, may be from 100% propellant and 0% solvent to 85% propellant and 15 % solvent. Within this range of percentages, pharmaceutically useful concentrations of Δ^9 THC can be achieved and droplet size is still small enough ($<5.8\mu\text{m}$) to provide excellent aerosol delivery of the drug. While these ratios reflect preferred embodiments of the invention, it will be recognized by those of skill in the art that the exact ratio of propellant to solvent (e.g. ethanol) may vary according to the desired final concentration of Δ^9 THC and droplet size. Any ratio of propellant to solvent that results in appropriate sized droplets and adequate dissolution of the Δ^9 THC may be used in the practice of this invention, and this will generally be in the range of from 100 to 80% propellant and 0 to 20% solvent. It is expected that a wide variety of solvents, such as ethanol, propanol, propylene glycol, glycerol, polyethylene glycol, etc. may be used in the preparation of formulations contemplated by this invention.

Those skilled in the art will also recognize that the "respirable dose" (or mass of Δ^9 THC in particles with aerodynamic diameters small enough to be delivered to and absorbed by the lungs) (Figure 1) may be increased by choosing MDI spray nozzles of different design and smaller orifice diameters. Respirable doses may also be increased by extending the mouthpiece of the MDI in such a way as to create an integral or separate aerosol spacer or reservoir attached to the mouthpiece of the MDI. This promotes an increase in droplet evaporation and hence in the percentage of the dose in smaller "respirable" particles or droplets. Generally, the optimal size of a respirable droplet is less than $10\mu\text{m}$ in size.

TABLE 3. Apparent Solubility of Δ^9 THC in Ethanol/HFA Propellant Blends				
Formulation	Mass (g) of Δ^9 THC in Sample	Mass (g) of Formulation Sampled	Apparent Solubility Mean (\pmSD)	Comments
Δ^9 THC in 100% HFA 134a	0.000240	0.1071	0.224% w/w (\pm 0.063)	Excess Δ^9 THC added to propellant blend (in pressurized MDI). Solubility sample removed using puff absorber. n=5
Δ^9 THC in 5% Ethanol / 95% HFA 134a	0.00144	0.0914	1.585% w/w (\pm 0.321)	As above
Δ^9 THC in 10% Ethanol / 90% HFA 134a	0.00363	0.1036	3.511% w/w (\pm 0.249)	As above
Δ^9 THC in 15% Ethanol / 85% HFA 134a	0.00536	0.1098	4.883% w/w (\pm 0.224)	As above
Δ^9 THC in 100% HFA 227	0.00021	0.1451	0.147% w/w (\pm 0.008)	As above
Δ^9 THC in 5% Ethanol / 95% HFA 227	0.00134	0.0979	1.339% w/w (\pm 0.169)	As above
Δ^9 THC in 10% Ethanol / 90% HFA 227	0.00454	0.1267	3.240% w/w (\pm 0.161)	As above
Δ^9 THC in 15% Ethanol / 85% HFA 227	0.00623	0.1062	5.940% w/w (\pm 0.191)	As above

A distinct advantage of the present formulations is that, surprisingly, the use of surface active agents or "surfactants" as valve lubricants and solubilizers is not necessary. This is in contrast to the invention of Purewal and Greenleaf (European Patent 0,372,777; reference #25) which provides HFA 134a/ethanol mixtures to produce stable formulations of pharmaceuticals in the presence of lipophilic surface active agents. Lipophilic surface active agents are incorporated in that invention in order to suspend undissolved material and to ensure adequate valve lubrication of the MDI. Without adequate valve lubrication, the useful life of the MDI and its ability to deliver an accurate dose of drug are severely attenuated. However, probably due to the inherent lubricity of the formulations of the present invention, the use of such surface active agents is unnecessary. This simplifies the composition and thus is an advantage with respect to cost and the elimination of potentially deleterious interactions between components of the formulations and the agents.

A major consideration in the formulation of any drug is its stability. Δ^9 THC is known to deteriorate upon storage so that the effective concentration decreases and the purity is vitiated. The stability of the formulations of the present invention were tested according to accelerated storage testing protocols. The results are given in Figure 1 and Tables 4A and 4B. The formulations of the present invention were shown to be stable with respect to the release of aerosolized Δ^9 THC in reproducible doses following accelerated storage testing. Apparently, the containment of Δ^9 THC in solution in the non-aqueous formulations of the present invention is excellent with respect to chemical degradation, making possible the construction of a multidose inhaler with a good shelf life prognosis.

Further, lipophilic materials like Δ^9 THC are generally known to partition into the elastomers of the valves in MDI formulations. (Δ^9 THC is highly lipophilic as reflected in its octanol:water partition coefficient of 6000:1). Over time, this partitioning results in a decrease in the emitted or delivered dose of a lipophilic drug. Thus, this phenomenon also decreases the useful shelf-life of such preparations. However, the data presented in Figure 1 and Table 4 show that this is not the case

with the formulations of the present invention. The emitted or delivered doses were constant over the time period tested. This may be due to the somewhat surprising preference of Δ^9 THC for the formulation itself, rather than for the valve elastomers.

TABLE 4A. Formulation and aerosol characteristics of Δ^9 THC pressurized metered dose inhalers in ethanol/hydrofluoroalkane (HFA) propellant blends.

Inhaler	Formulation (% w/w)			Description
	Δ^9 THC	Ethanol	Propellant	
1	0.13%	~5%	95% HFA 134a	3/98 Pale Yellow Solution
2	0.13%	~5%	95% HFA 227	3/98 Pale Yellow Solution
3	0.12%	~5%	95% HFA 134a	3/98 Pale Yellow Solution
4	0.18%	~5%	95% HFA 134a	3/98 Pale Yellow Solution
5	0.27%	~5%	95% HFA 227	3/98 Pale Yellow Solution
6	0.25%	~5%	95% HFA 134a	3/98 Pale Yellow Solution
7	0.57%	~5%	95% HFA 134a	3/98 Yellow Solution
8	0.58%	~5%	95% HFA 227	3/98 Yellow Solution
9	0.49%	~5%	95% HFA 134a	3/98 Yellow Solution
10	1.02%	~5%	95% HFA 134a	3/98 Yellow Solution
11	1.11%	~5%	95% HFA 227	3/98 Yellow Solution
12	0.97%	~5%	95% HFA 134a	3/98 Yellow Solution
SS* #1 Initial	1.07%	4.94%	94.0% HFA 134a	6/98 Yellow Solution

SS* #1 after 28 days at 40°C/82% RH**	1.07%	4.94%	94.0% HFA 134a	7/98 Yellow Solution
SS* #2 after 21 days at 40°C/82% RH**	1.00%	5.01%	95% HFA 134a	7/98 Yellow Solution
SS* #3 Modified Actuator** *	1.02%	5.15%	93.8% HFA 134a	10/98 Yellow Solution

^a Mean (Standard Deviation) of five determinations.

^b Mass of Δ^9 THC aerosol particles <5.8 μm aerodynamic diameter

*SS: Stability Sample

** RH: relative humidity

*** Approximate spray nozzle diameter = 0.2 mm.

TABLE 4B: Formulation and aerosol characteristics of Δ^9 THC pressurized metered dose inhalers in ethanol/hydrofluoroalkane (HFA) propellant blends.

Inhaler	Aerosol Characterization		
	Metered Dose (mg) ^a	Emitted Dose (mg) ^a	Fine Particle Dose (mg) _{a,b}
11	1.72 (0.25)	1.32 (0.17)	ND
12	0.94 (0.23)	0.97 (0.10)	0.38 (0.02)

SS* #1 Initial	1.10 (0.07)	0.90 (0.03)	0.22 (0.03)
SS* #1 after 28 days at 40°C/82% RH**	1.06 (0.03)	0.92 (0.04)	0.23 (0.02)
SS* #2 after 21 days at 40°C/82% RH**	1.02 (0.05)	0.90 (0.05)	0.21 (0.02)
SS* #3 Modified Actuator***	ND	ND	0.40 (n=1)

^a Mean (Standard Deviation) of five determinations.

^b Mass of Δ^9 THC aerosol particles with $<5.8 \mu\text{m}$ aerodynamic diameter

*SS: Stability Sample

** RH: relative humidity

ND: not
determined

*** Approximate spray nozzle diameter = 0.2 mm

The final concentration of Δ^9 THC in a given formulation may be varied by adjusting the ratio of propellant to solvent and thus the solubility of the Δ^9 THC. Higher percentages of solvent (e.g. ethanol) generally allow a higher amount of Δ^9 THC to be dissolved. For example, in preferred embodiments of the invention, the apparent solubility of Δ^9 THC ranged from 0.147% w/w to 5.94% w/w as the propellant composition varied from 100% HFA 227 to 85% HFA 227 and 15% ethanol. Thus, the dose of Δ^9 THC in a given metered volume may be selected by changing the formulation.

Further, as stated above, the "fine particle dose" or "respirable dose" of a drug dispensed with an MDI is a function of the spray nozzle diameter. In Figure 1 and Tables 4A and 4B, the spray nozzle diameter is 0.4mm. The "fine particle dose" or "respirable dose" of the formulations of the present invention was shown to be

unaffected by storage.

The Δ^9 THC of the present invention is pharmaceutically pure. That is, its form is the nonionized resinous drug substance (6aR-trans)-6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-6H-dibenzo[b,d]-pyran-1-ol. Although its preferred embodiment in this invention is not a salt or ester, it will be readily understood by those of skill in the art that other appropriate forms of Δ^9 THC may be synthesized (e.g. esters and salts) and thus used in the practice of this invention.

The desired final concentration of Δ^9 THC in a patient's serum will vary from patient to patient depending on, for example, the nature and severity of the condition being treated, and the patient's overall condition, weight, gender and response to the drug, etc. But the desired range will generally be 10-100ng/ml at 15 minutes following inhalation. The level of Δ^9 THC in a patient's serum can be readily and reliably monitored by gas chromatography/mass spectrophotometry (GC/MS).

The exact treatment protocol to be used may vary from patient to patient depending on the circumstances. For example, in a preferred embodiment of the invention, a patient receiving chemotherapy may have one dose of Δ^9 THC prescribed via inhalation, to be administered 15 minutes before chemotherapy and 4-8 times daily following chemotherapy. In another preferred embodiment, a patient suffering from anorexia associated with AIDS wasting syndrome may have Δ^9 THC by inhalation prescribed 3-5 times daily, 30 minutes before each meal or snack. In other preferred embodiments, a patient suffering from cancer pain, or spasticity related to either multiple sclerosis or spinal cord injury may have Δ^9 THC by inhalation prescribed 3-6 times daily. Those skilled in the art will readily recognize that the treatment protocol may be crafted so as to address the particular needs of each individual patient on a case by case basis.

Δ^9 THC may be used alone or in combination with other medications. Those skilled in the art will readily recognize that, for example, in the case of AIDS wasting syndrome, the patient will likely also be taking drugs that combat the AIDS virus. Similarly, those skilled in the art will readily recognize that patients receiving

chemotherapy for cancer may also receive other antiemetics, and cancer patients seeking to relieve pain are likely to receive opioids as well as nonsteroidal anti-inflammatory agents.

The containers for the formulations of the instant invention may be any that are suitable for the efficacious delivery of aerosol inhalants. Several containers and their method of usage are known to those of skill in the art. For example, MDIs can be used with various dose metering chambers, various plastic actuators and mouthpieces, and various aerosol holding chambers (e.g. spacer and reservoir devices), so that appropriate doses of Δ^9 THC reach and deposit in the lung and are thereafter absorbed into the bloodstream. In addition, a lock mechanism such as that shown in U.S. Patent 5,284,133 to Burns and Marshak, which is herein incorporated by reference, can be used to prevent overdose or unauthorized consumption of Δ^9 THC. Figure 2 provides a generalized drawing of an MDI containing the composition of this invention and provides the advantage of delivering metered quantities of Δ^9 THC on a repetitive basis. The MDI includes a container 100 for holding the composition and a valve delivery mechanism 102 for delivery of aerosolized Δ^9 THC.

While the invention has been described in terms of its preferred embodiments, those skilled in the art will recognize that the invention can be practiced with modification within the spirit and scope of the appended claims.

REFERENCES

1. Workshop on the medical utility of marijuana. National Institutes of Health, August 1997.
2. Beal, J.A., Olson, R., Lefkowitz, L., Laubenstein, L., Bellman, P., Yangco, B., Morales, J.O., Murphy, R., Powderly, W., Plasse, T.F., Mosdell, K.W. and Shepard, K.W. (1997) Long-term efficacy and safety of dronabinol for acquired immunodeficiency syndrome-associated anorexia. *J. Pain. Symptom Manage.* 14:7-14.

3. Beal, J.A., Olson, R., Laubenstein, L., Morales, J.O., Bellman, B., Yangco, B., Lefkowitz, L., Plasse, T.F. and Shepard, K.V. (1995) Dronabinol as a treatment for anorexia associated with weight loss in patients with AIDS *J. Pain. Symptom Manage.* 10:89-97.
4. McCabe, M., Smith, F.P., MacDonald, J.S., Wooley, P.V., Goldberg, D. and Schein, P.S. (1988) Efficacy of tetrahydrocannabinol in patients refractory to standard antiemetic therapy. *Invest. New Drugs* 6:243-246.
5. Lucas, V.S. and Laszlo, J. (1980) Δ^9 -THC for refractory vomiting induced by cancer chemotherapy. *JAMA* 243:1241-1243.
6. Sallan, S.E., Cronin, C., Zelen, M. and Zinberg, N.E. (1980) Antiemetics in patients receiving chemotherapy for cancer: a randomized comparison of Δ^9 THC and prochlorperazine. *N. Engl. J. Med.* 302:135-138.
7. Frytak, S., Moertel, C.G., O'Fallon, J.R., Rubin, J., Creagan, E.T., O'Connell, M.J., Schutt, A.J. and Schwartz, N.W. (1979) Delta-9-tetrahydrocannabinol as an antiemetic for patients receiving cancer chemotherapy: a comparison with prochlorperazine and a placebo. *Ann. Inter. Med.* 91:825-830.
8. Chang, A.E., Shilling, D.J., Stillman, R.C., Goldgerg, N.H., Seipp, C.A., Barofdky, I., Simon, R.M. and Rosenberg SA (1979) Δ^9 THC as an antiemetic in cancer patients receiving high-dose methotrexate. *Ann. Internal. Med.* 91:819-824.
9. Sallan, S.E., Zinberg, N.E. and Frei, I.E. (1975) Antiemetic effect of Δ^9 THC in patients receiving cancer chemotherapy. *New Engl. J. Med.* 293:795-797.
10. Noyes, J.R., Brunk, S.F., Baram, D.A. and Canter, A. (1975) The analgesic

properties of Δ^9 THC and codeine. *J. Clin. Pharmacol.* 15:139-143.

11. Noyes, R., Jr., Brunk, S.F., Baram, D.A. and Canter, A. (1975) Analgesic effect of Δ^9 -tetrahydrocannabinol. *J. Clin. Pharmacol.* 15:139-143.
12. Brenneisen, R., Egli, A., Elosohly, M.A., Henn, V. and Spiess, Y. (1996) The effect of orally and rectally administered Δ^9 THC on spasticity: a pilot study with 2 patients. *Int. J. Clin. J. Pharmacol. Ther.* 34:446-452.
13. Ungerleider, J.T., Andyrsiak, T.F.L., Ellison, G.W. and Myers, L.W. (1987) Δ^9 THC in the treatment of spasticity associated with multiple sclerosis. *Adv. Alcohol Subst. Abuse* 7:39-50.
14. Clifford, D.B. (1983) Tetrahydrocannabinol for tremor in multiple sclerosis. *Ann. Neurol.* 13:669-171.
15. Petro, D.J. and Ellenberger, C. (1981) Treatment of human spasticity with delta 9-tetrahydrocannabinol. *J. Clin. Pharmacol.* 21:413S-416S.
16. Maurer, M., Henn, V., Dittrich, A. and Hofman, A. (1990) Delta 9-tetrahydrocannabinol shows antispastic and analgesic effects in a single case double-blind trial. *Eur. Arch. Psychiatry Neurol. Sci.* 240:1-4.
17. Merritt, J., Crawford, W., Alexander, P., Anduze, A. and Gelbart, S. (1980) Effects of marihuana on intra ocular and blood pressure in glaucoma. *Opht.* 87:222-228.
18. Cooler, P. and Gregg, J.M. (1977) Effect of delta 9- Δ^9 THC on intra ocular pressure in humans. *South. Med. J.* 70:951-954.

19. PDR (1995) Physician's Desk Reference (49) Montvalek, New Jersey: Medical Economics Data Production Co., pp.2787.
20. Ohlsson, A., Lindgren, J.E., Wahlen, A., Agurall, S., Hollister, L.E. and Gillespie, H.K. (1980) Plasma Δ^9 THC concentrations and effects after oral and intravenous administration and smoking. *Clin. Pharmacol. Ther.* 28:409-416.
21. Olsen, J.L., Lodge, J.W., Shapiro, B.J. and Tashkin, D.P. (1976) An inhalation aerosol of Δ^9 - tetrahydrocannabinol. *J. Pharmacy and Pharmacol.* 28:86.
22. Dalby, R.N. and Byron, P.R. (1988) Comparison of output particle size distributions from pressurized aerosols formulated as solutions or suspensions. *Pharm. Res.* 5:36-39.
23. Tashkin, D.P., Reiss, S., Shapiro, B.J., Calvarese, B., Olsen, J.L. and Lidgek, J.W. (1977) Bronchial effects of aerosolized Δ^9 - tetrahydrocannabinol in healthy and asthmatic subjects. *Amer. Rev. of Resp. Disease.* 115:57-65.
24. Williams, S.J., Hartley, J.P.R. and Graham, J.D.P. (1976) Bronchodilator effect of delta-9-THC administered by aerosol to asthmatic patients. *Thorax.* 31:720-723.
25. European Patent 0,372,777 (Riker Laboratories). Medicinal aerosol formulations.

CLAIMS

We claim:

- 1 1. An aerosol-dispensable pharmaceutical composition, comprising:
2 a hydrofluoroalkane propellant; and
3 Δ^9 - tetrahydrocannabinol present in a pharmaceutically acceptable form
4 and at a
5 pharmaceutically effective concentration dissolved in said hydrofluoroalkane
6 propellant.
- 1 2. The aerosol dispensable pharmaceutical composition of claim 1 wherein said
2 Δ^9 - tetrahydrocannabinol is dissolved in said hydrofluoroalkane propellant without the
3 use of a solubilizing agent selected from the group consisting of solvents and
4 surfactants.
- 1 3. The aerosol-dispensable pharmaceutical composition of claim 1 wherein said
2 hydrofluoroalkane is selected from the group consisting of: HFA 134a and HFA 227.
- 1 4. The aerosol-dispensable pharmaceutical composition of claim 3 wherein said
2 hydrofluoroalkane is present in an amount in excess of 85% by weight.
- 1 5. The aerosol-dispensable pharmaceutical composition of claim 1 further
2 comprising an organic solvent.
- 1 6. The aerosol-dispensable pharmaceutical composition of claim 5 wherein said
2 organic solvent is ethanol present in an amount ranging up to 15% by weight.
- 1 7. The aerosol-dispensable pharmaceutical composition of claim 1 wherein said
2 pharmaceutically effective concentration of Δ^9 -tetrahydrocannabinol ranges from 0.05
3 to 10% by weight.

1
2 8. The aerosol-dispensable pharmaceutical composition of claim 1 wherein said
3 pharmaceutically effective concentration of Δ^9 -tetrahydrocannabinol ranges from 0.1
4 to 6% by weight.

1 9. The aerosol-dispensable pharmaceutical composition of claim 1 wherein Δ^9 -
2 tetrahydrocannabinol is present in pure form.

1 10. A solvent free pharmaceutical composition consisting essentially of
2 1,1,1,2,3,3,3-hydrofluoropropane (HFA 227) and Δ^9 - tetrahydrocannabinol.

1 11. A solvent free pharmaceutical composition consisting essentially of
2 1,1,1,2-hydrofluroethane (HFA 134a) and Δ^9 - tetrahydrocannabinol.

1 12. A method of treating a patient in need thereof with an aerosolized
2 pharmaceutically acceptable form of Δ^9 - tetrahydrocannabinol, comprising the step of:
3 administering an aerosolized dose of a pharmaceutically acceptable form of
4 Δ^9 -tetrahydrocannabinol as respirable droplets to a patient's lung from a composition
5 comprised of a hydrofluoroalkane propellant and said pharmaceutically acceptable
6 form of Δ^9 - tetrahydrocannabinol.

1 13. The method of claim 12 wherein said composition comprises a
2 pharmaceutically acceptable solvent.

3 14. The method of claim 12 wherein said composition is solvent-free.

4 15. The method of claim 12 wherein said aerosolized dose is sufficient to
5 reduce nausea.

1 16. The method of claim 12 wherein said aerosolized dose is sufficient to
2 reduce vomiting.

1 17. The method of claim 12 wherein said aerosolized dose is sufficient to
2 reduce pain.

1 18. The method of claim 12 wherein said aerosolized dose is sufficient to
2 relieve muscle spasticity.

1 19. The method of claim 12 wherein said aerosolized dose is sufficient to
2 relieve migraine headaches.

1 20. The method of claim 12 wherein said aerosolized dose is sufficient to
2 relieve movement disorders.

1 21. The method of claim 12 wherein said aerosolized dose is sufficient to
2 increase appetite in patients suffering from cachexia.

1 22. The method of claim 12 wherein said pharmaceutically acceptable form of
2 Δ^9 - tetrahydrocannabinol is pure Δ^9 - tetrahydrocannabinol and said hydrofluoroalkane
3 is selected from the group consisting of HFA 134a and HFA 227.

4 23. A metered dose inhaler, comprising,
5 a housing;
6 a metering valve connected to said housing; and,
7 an aerosol-dispensable pharmaceutical composition which includes a
8 hydrofluoroalkane propellant and Δ^9 - tetrahydrocannabinol present in a
9 pharmaceutically effective concentration dissolved in said hydrofluoroalkane
10 propellant.




FIG.1





FIG.2



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 31/35, A61L 9/04, A61M 11/00, A01N 43/16	A3	(11) International Publication Number: WO 00/24362 (43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/24486 (22) International Filing Date: 20 October 1999 (20.10.99) (30) Priority Data: <div style="display: flex; justify-content: space-between;"> <div>60/105,850</div> <div>27 October 1998 (27.10.98)</div> <div>US</div> </div> <div style="display: flex; justify-content: space-between;"> <div>09/273,766</div> <div>22 March 1999 (22.03.99)</div> <div>US</div> </div> (71) Applicant: VIRGINIA COMMONWEALTH UNIVERSITY [US/US]; 1101 E. Marshall Street, Richmond, VA 23298 (US). (72) Inventors: PEART, Joanne; Apartment #1332, 11808 Chase Wellesley Drive, Richmond, VA 23233 (US). BYRON, Peter; 8101 Lower Ralston Court, Richmond, VA 23229 (US). LICHTMAN, Aron; 2306 Stonemill Court, Richmond, VA 23233 (US). MARTIN, Billy; 2406 St. Regis Drive, Richmond, VA 23233 (US). (74) Agent: WHITMAN, Michael, E.; Whitham, Curtis & Whitham, Suite 900, 11800 Sunrise Valley Drive, Reston, VA 20191 (US).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 24 August 2000 (24.08.00)	
(54) Title: Δ^9 TETRAHYDROCANNABINOL (Δ^9 THC) SOLUTION METERED DOSE INHALERS AND METHODS OF USE (57) Abstract <p>The present invention provides therapeutic formulations for solutions of Δ^9-tetrahydrocannabinol to be delivered by metered dose inhalers (100), (102). The formulations, which utilize non-CFC propellants, provide a stable aerosol-deliverable source of Δ^9-tetrahydrocannabinol for the treatment of various medical conditions, such as nausea, and vomiting associated with chemotherapy; muscle spasticity; pain; anorexia associated with AIDS wasting syndrome; epilepsy; glaucoma; bronchial asthma; and mood disorders.</p>		
		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24486

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/35; A61L 9/04; A61M 11/00; A01N 43/16;

US CL : 128/200.14, 200.23, 203.12; 424/45; 514/455

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 128/200.14, 200.23, 203.12; 424/45; 514/455

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	U.S. 4,635,651 A (JACOBS) 13 January 1987, Col. 2 lines 25-33.	1-23
Y	U.S. 5,502,076 A (DIXIT et al.) 26 March 1996, col. 1 lines 23--27, and col. 2 lines 60-63.	3, 4, 10, 11, 12, 22, 23
Y	U.S. 5,804,592 A (VOLICER) 08 September 1998, col. 2 lines 1, 2, 6-9, 21, 22, 34, 41-42; and col. 4 line 40.	15-21

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 MARCH 2000

Date of mailing of the international search report

02 MAY 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230

Authorized officer

AARON J. LEWIS

Telephone No. (703) 308-0716

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24486

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☒

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24486

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-11, drawn to a pharmaceutical composition.

Group II, claims 12-23, drawn to method, and apparatus for administering an aerosolized dose of delta-9-tetrahydrocannabinol to a patient's lung.

Groups I and II, the inventions listed as these groups do not relate to a single inventive concept under PCT Rule 13.1 because under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The process, the apparatus as claimed can be practiced with another materially different product, such as lung surfactants, insulin, nitric oxide, and anesthetics.

